



CORNELL COOPERATIVE EXTENSION - SUFFOLK COUNTY



INSECT AND PLANT DISEASE DIAGNOSTIC LABORATORY

EDUCATION CENTER
423 GRIFFING AVENUE
RIVERHEAD, NY 11901
HORT INFO LINE 631.727.4126



Cornell University
Cooperative Extension
of Suffolk County

BAYARD CUTTING ARBORETUM
MONTAUK HWY. PO BOX 463
OAKDALE, NY 11769
HORT INFO LINE 631.581.4223

Summer Patch on Turfgrass *Magnaporthe poae*



Figure 1. Typical “patch” symptom of summer patch disease. (North Carolina State University)



Figure 2. A lawn exhibiting symptoms of summer patch infection. (Note that these symptoms can resemble those associated with other common turfgrass problems.) (T. Turner, HGIC, University of Maryland)

Introduction: Kentucky bluegrass (*Poa pratensis*), annual bluegrass (*Poa annua*), and fescues (*Festuca* sp.) can be affected during the summer by an interaction of environmental factors and a root and/or crown rot caused by the fungus *Magnaporthe poae*. This disease is known as Summer Patch. Bentgrasses (*Agrostis* sp.) may also become infected but show few symptoms and may continue to perform where other grasses decline.

Summer Patch occurs between June and September. This disease, in combination with another disease now known as Necrotic Ring Spot were once grouped together into the disease complex previously known as Fusarium Blight. The high level of difficulty involved in clinical evaluations of *Leptosphaeria korrae*, *Magnaporthe poae*, and related fungi, and the extreme ease of isolation of decomposer fungi such as *Fusarium* species, originally led to the misnaming. Eventually, these organisms and their symptoms were separated out into the diseases now known as [Necrotic Ring Spot](#) and Summer Patch. The latter will be described in greater detail here.

Symptoms: It can be difficult to diagnose this disease by symptoms alone in the early stages. The disease begins as scattered small round patches of thin, wilted or slow growing turf. Initially, affected patches (**Fig. 1**) may be only 3-8 cm in diameter, but they may enlarge to about 30 cm in diameter (about 12 inches) and range from gray-green to light tan or straw-colored (**Fig. 2**). In rarer circumstances, they may get to be twice that size. As patches enlarge, they may coalesce, and form

crenscents of yellow or tan turf. Where turf within the center of a patch begins to recover, necrotic rings may




Figure 3. Crown and root rotting symptoms on creeping bentgrass. (J. P. Kerns, Plant Pathology, North Carolina State University)

become evident. As Necrotic Ring Spot may exhibit similar symptoms, microscopic examination is often necessary to determine the cause of the problem. The roots, crowns and stolons of infected plants may appear to be dark brown (**Fig. 3**) as the dark mycelium of the fungus (**Fig. 4**) invades the tissue. As the disease progresses, the cortex may begin to rot and plants may die.

Summer Patch blighted areas often occur on lawn sites that receive direct sun and are on south-facing slopes, or near sidewalks, driveways, buildings, or other "hot spots" or otherwise stressed areas in the yard or on golf courses. In the cool weather of autumn, the grass may begin to grow into these dead areas again. The disease, however, is likely to reappear in previously affected areas the following summer, and to increase in intensity.

Summer patch usually occurs during the hotter periods (June, July, and August) of the year. Summer patch is less of a problem during cool summers with adequate rainfall.

Disease Cycle: The fungus, *Magnaporthe poae*, survives unfavorable conditions as mycelium in infected plant tissue or plant debris. The optimum temperature for fungal growth is 28°C. Infection takes place in late spring when soil temperatures reach 19-20°C. Spread during the growing season occurs as the fungus grows between roots. Symptoms may not be evident until the temperature increases very drastically during wet weather. The pathogen may also be spread by movement of infected plant material and on mechanical equipment.

Management Strategies: The primary stresses that influence disease development include excesses of thatch, fertilizer, and turf canopy temperature, as well as incorrect timing of fertilizer applications, low mowing height, and pH extremes. Each of these stresses can be




Figure 4. A microscopic image of the dark runner hyphae of the fungus *Magnaporthe poae*. (J.P. Kerns, Plant Pathology, North Carolina State University)

reduced through appropriate cultural practices described below.

Disease severity may worsen at a higher pH, so try to maintain the pH of the soil and Rhizosphere at 5.5 to 6.0. Use an acidifying fertilizer where the pH is above 6.0, and try to avoid the use of products that may raise the pH. Contact Cornell Cooperative Extension – Suffolk County [for information on having soil tested for pH](#).

For information on fertilizing Kentucky bluegrass lawns on Long Island contact Cornell Cooperative Extension – Suffolk County. Do not apply even small amounts of fertilizer during the June-August stress period because this will tend to stimulate the disease.

Deep watering is essential for proper root growth. Water the soil under disease-prone areas to a depth of 15 to 20 cm every 7-10 days during the dry periods of the summer. Soaker hoses are very useful for supplementary watering on steeper slopes where other sprinklers are inefficient. The harmful effects of excessive temperature can be reduced by a light sprinkling of the surface at mid-day.

Proneness to disease in turf is increased as the cutting height is decreased. Cut lawns at 5 to 10 cm height, and often enough so that less than 1/3 of the leaf blade is removed during each mowing.

Thatch (the layer of organic matter between the mineral soil and the green grass) should be no more than 1.5 cm (1/2 inches) in thickness. Thatch can be removed by vertical slicing machines and/or aeration during the spring and early fall. Over a longer period thatch will be reduced by using the cultural practices discussed above.

Kentucky bluegrass cultivars such as Adelphi, America, Aspen, Columbia, Eclipse, Glade, Midnight, Nassau, Parade, Ram I, Sydsport, Touchdown, Vantage, Windsor, and Victa are less susceptible to summer patch than others. Blend seed of resistant cultivars with that of one or more otherwise desirable cultivars. Blending 10-15% (by weight) of perennial ryegrass seed into bluegrass seed will prevent this disease from occurring. Ryegrass can also be seeded into existing lawns.

Chemical treatment is efficient only when the previously mentioned cultural practices are first used. Furthermore, applications must be made before the crown rot develops sufficiently to cause visual symptoms of the disease. Fungicides containing the active ingredients azoxystrobin, cyproconazole, fenarimol, propiconazole, iprodione, thiophanate-methyl, trifloxystrobin or triadimefon are available for control of summer patch in commercial applications. Other products containing azoxystrobin, myclobutanil, thiophanate-methyl, or triadimefon may be available for use on home lawns by homeowners to manage this disease.

Thoroughly water (applying 2 to 3 cm of water) areas with a history of this disease several days before applying the fungicide. To determine the best time to treat, monitor soil temperature to a depth of 2 inches, and make the first application when the soil temperature reaches 55°F. or about 13°C. In upstate New York, the first application may be made in early to mid-June with a second application two to three weeks later. In southeastern New York or on Long Island, the first application needs to be made a few to several weeks earlier (mid-April) depending on the soil temperature. Additional applications may be required for certain fungicides. Always adhere to the rates and procedures recommended on the fungicide package label.

Reprinted from: *Summer Patch on Turfgrass Magnaporthe poae*, The Plant Disease Diagnostic Clinic at Cornell University, Ithaca, NY; Updated, SLJ, 11/07.

Slight revision T. Kowalsick & T. Yeh, 1/2008

The Pesticide Management Education Program (PMEP), in cooperation with the New York State Department of Environmental Conservation (NYSDEC), maintains a web site with a searchable database for pesticide products currently registered in New York State. Homeowners who have Internet access can locate currently registered products at <http://pmp.cce.cornell.edu/pims/current>. Several different queries are available that will produce a summary for the product(s) that the system locates. If the system fails to locate the product in question, then that product is not currently registered in New York State. The database also provides a summary of important information related to every product currently registered. Two data fields "Status" and "Expiration Date" are provided in each summary. Products with a status of "Registered - Discontinued" are currently registered but will probably be discontinued for use, sale, and distribution in New York State after the date noted in the "Expiration Date" field.

This publication contains pesticide recommendations. Changes in pesticide regulations occur constantly, some materials mentioned may no longer be available, and some uses may no longer be legal. All pesticides distributed, sold, and/or applied in New York State must be registered with the New York State Department of Environmental Conservation (DEC). Questions concerning the legality and/or registration status for pesticide use in New York State should be directed to the appropriate Cornell Cooperative Extension Specialist or your regional DEC office. READ THE LABEL BEFORE APPLYING ANY PESTICIDE.

TK: 1/2008